

## Lipidomics study of mesenchymal stromal cells derived from human placenta

Martina Rossi<sup>1</sup>, Alexandros Chatgillialoglu<sup>2</sup>, Francesco Alviano<sup>1</sup>, Cosetta Marchionni<sup>1</sup>, Paola Poggi<sup>2</sup>, Francesca Ricci<sup>3</sup>, Pier Luigi Tazzari<sup>3</sup>, Elisabetta Caramelli<sup>1</sup>, Laura Bonsi<sup>1</sup>

<sup>1</sup> Department of Experimental, Diagnostic and Specialty Medicine, University of Bologna, Bologna, Italy -

<sup>2</sup> Remembrance Srl, Imola, Italy - <sup>3</sup> Service of Immunohematology and Transfusion Medicine, S. Orsola University Hospital, Bologna, Italy

The interest for lipid metabolism in the stem cell field has increased in the last few years (1,2,3). Membrane lipidomics embraces many aspects of cell metabolism and the role of lipids is now considered more than merely inert and structural in delimitating the extra- and intra-cellular compartments (4,5). Nevertheless, we are still far from understanding the impact of membrane lipidomics in stemness maintenance and differentiation patterns.

The aim of our work was to study membrane lipidomics of mesenchymal stromal cells derived from human placenta and correlate it to specific biological properties, by using chemically-defined tailored lipid supplements (Refeed®). In the experimental study, the cell membranes of freshly isolated mesenchymal stromal cells obtained from human fetal membranes (FM-MSCs) were characterized for fatty acid composition. Then, we investigated cell morphology, viability, proliferation, differentiation and immunomodulation after in-vitro exposure to Refeed® supplements. Control MSCs were cultured without lipid supplementation.

Our results showed a significant reduction of membrane fluidity for in-vitro primary cells, with cell membrane fatty acid composition greatly differing from the in-vivo one. By tailoring lipid supplementation, the fatty acid composition and biophysical properties of in-vitro cell membranes resulted more similar to the in-vivo counterparts, with higher omega-6 fatty acid content and increased membrane fluidity. These modifications of membrane composition and properties had no effect on cell morphology and viability, whereas ameliorated cell proliferation rate, differentiation ability and immunomodulatory properties. In particular, supplemented FM-MSCs showed an increased expression of cell membrane molecules like Vascular Endothelial Growth Factor Receptors 1 (VEGFR-1 or Flt-1) and 2 (VEGFR-2 or KDR), that correlated with a more efficient response to angiogenic commitment. Moreover, regarding immunomodulation, supplemented FM-MSCs displayed an increased expression of the tolerogenic cell surface protein HLA-G, that positively influenced the in-vitro cell immunomodulatory ability.

Finally, these data suggest that specific lipid supplementation have functional consequences on in-vitro MSC behavior and may influence cell-based therapeutic approaches.

### References

- [1] Yanes O, Clark J, Wong DM *et al*, Metabolic oxidation regulates embryonic stem cell differentiation. *Nat Chem Biol* 2010; 6(6):411-7.
- [2] Ito K, Carracedo A, Weiss D *et al*, A PML-PPAR- $\delta$  pathway for fatty acid oxidation regulates hematopoietic stem cell maintenance. *Nat Med* 2012; 18(9):1350-8.
- [3] Knobloch M, Braun SM, Zurkirchen L *et al*, Metabolic control of adult neural stem cell activity by Fasn-dependent lipogenesis. *Nature* 2013; 493(7431):226-30.
- [4] Langelier B, Linard A, Bordat C *et al*, Long chain-polyunsaturated fatty acids modulate membrane phospholipid composition and protein localization in lipid rafts of neural stem cell cultures. *J Cell Biochem* 2010; 110(6):1356-64.
- [5] Smith AN, Muffley LA, Bell AN *et al*, Unsaturated fatty acids induce mesenchymal stem cells to increase secretion of angiogenic mediators. *J Cell Physiol* 2012; 227(9):3225-33.

### Keywords

Mesenchymal stromal cells; cell membranes; lipid metabolism; fatty acids.